

NOMENCLATORIAL NOTES XIII

BY

CHARLES SCHWEINFURTH

Erythrodes santensis (*Kränzl.*) *C. Schweinfurth*
comb. nov.

Physurus santensis Kränzlin in Kungl. Svensk. Vet.
Akad. Handl. 46 (1911) 39, t. 7, fig. 6.

Since *Physurus* L.C. Rich., a nomen nudum, is synonymous with *Erythrodes* Bl., the new combination is necessary.

It is evident that the artist's drawing (l.c.) represents the natural forcipate position of the apical arms of the lip, whereas in the forcibly expanded position these arms are reversed and assume a retrorsely lunate position.

A collection from Colombia appears to be referable to this species which was described from São Paulo, Brazil. These plants differ from the typical form in having broader leaves (up to about 20 mm. wide) and in having larger petals.

COLOMBIA: Intendencia El Chocó, in dense forest south of Río Condoto, between Quebrada Guarapo and Mandinga, at 120-180 meters altitude, April 22, 28, 1939, *E.P. Killip 35674*.

Malaxis liparidioides (*Finet*) *C. Schweinfurth*
comb. nov.

Microstylis liparidioides Finet in Bull. Soc. Bot. France
55 (1908) 333, t. 10, figs. 1-10

As the concept *Microstylis* Nutt. ex Lindl. is synonymous with the earlier name *Malaxis* Soland. ex Sw., the new combination is required.

This species, which was described from Ecuador and Peru, is now recorded from Colombia.

COLOMBIA: Dept. El Valle, El Silencio, Yanaconas, at 1900-2200 meters altitude, terrestrial in dense forest, flowers green with white lip, February 28, 1939, *E.P. Killip & Hernando Garcia 33775*.